

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 1/20, 1/00, G01N 33/48 G01N 33/566	A1	(11) International Publication Number: WO 92/05244 (43) International Publication Date: 2 April 1992 (02.04.92)
(21) International Application Number: PCT/US91/06605 (22) International Filing Date: 12 September 1991 (12.09.91) (30) Priority data: 581,714 13 September 1990 (13.09.90) US (71) Applicant: DUKE UNIVERSITY [US/US]; Erwin Road, Durham, NC 27706 (US). (72) Inventors: KING, Klim ; 3925 Linden Terrace, Durham, NC 27705 (US). DOHLMAN, Henrik, G. ; 2226 Stewart Street, Berkeley, CA 94705 (US). CARON, Marc, G. ; 2606 Evans Drive, Durham, NC 27705 (US). LEFKOWITZ, Robert, J. ; 3539 Hamstead Court, Durham, NC 27707 (US).		(74) Agents: SIBLEY, Kenneth, D. et al.; Bell, Seltzer, Park & Gibson, P.O. Drawer 34009, Charlotte, NC 28234 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: EXPRESSION OF G PROTEIN COUPLED RECEPTORS IN YEAST (57) Abstract Disclosed is a transformed yeast cell containing a first heterologous DNA sequence which codes for a mammalian G protein coupled receptor and a second heterologous DNA sequence which codes for a mammalian G protein α subunit (mammalian G_α). The first and second heterologous DNA sequences are capable of expression in the cell, but the cell is incapable of expressing an endogenous G protein α -subunit (yeast G_α). The cells are useful for screening compounds which affect the rate of dissociation of G_α from $G_{\beta\gamma}$ in a cell. Also disclosed is a novel DNA expression vector useful for making cells as described above. The vector contains a first segment comprising at least a fragment of the extreme amino-terminal coding sequence of a yeast G protein coupled receptor. A second segment is positioned downstream from the first segment (and in correct reading frame therewith), with the second segment comprising a DNA sequence encoding a heterologous G protein coupled receptor.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE ⁺	Germany	MC	Monaco	US	United States of America
DK	Denmark				

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

EXPRESSION OF G PROTEIN COUPLED RECEPTORS IN YEAST

This invention was made with government support under NIH grant HL16037. The government may have certain rights to this invention.

Field of the Invention

5 This invention relates to yeast cells expressing heterologous G protein coupled receptors, vectors useful for making such cells, and methods of using the same.

Background of the Invention

10 The actions of many extracellular signals (for example, neurotransmitters, hormones, odorants, light) are mediated by receptors with seven transmembrane domains (G protein coupled receptors) and heterotrimeric guanine nucleotide-binding regulatory
15 proteins (G proteins). See H. Dohlman, M. Caron, and R. Lefkowitz, Biochemistry 26, 2657 (1987); L. Stryer and H. Bourne, Ann. Rev. Cell Biol. 2, 391 (1986). Such G protein-mediated signaling systems have been identified in organisms as divergent as yeast and man. See H.
20 Dohlman et al., supra; L. Stryer and H. Bourne, supra; K. Blumer and J. Thorner, Annu. Rev. Physiol. (in press). The β 2-adrenergic receptor (β AR) is the prototype of the seven-transmembrane-segment class of

-2-

ligand binding receptors in mammalian cells. In response to epinephrine or norepinephrine, β AR activates a G protein, G_s , which in turn stimulates adenylate cyclase and cyclic adenosine monophosphate production in the cell. See H. Dohlman et al., supra; L. Stryer and H. Bourne, supra. G protein-coupled pheromone receptors in yeast control a developmental program that culminates in mating (fusion) of a and α haploid cell types to form the a/α diploid. See K. Blumer and J. Thorner, supra; I. Herskowitz, Microbiol. Rev. 52, 536 (1988).

The present invention is based on our continued research into the expression of heterologous G protein coupled receptors in yeast.

Summary of the Invention

A first aspect of the present invention is a transformed yeast cell containing a first heterologous DNA sequence which codes for a mammalian G protein coupled receptor and a second heterologous DNA sequence which codes for a mammalian G protein α subunit (mammalian G_α). The first and second heterologous DNA sequences are capable of expression in the cell, but the cell is incapable of expressing an endogenous G protein α -subunit (yeast G_α). The cell optionally contains a third heterologous DNA sequence, with the third heterologous DNA sequence comprising a pheromone-responsive promoter and an indicator gene positioned downstream from the pheromone-responsive promoter and operatively associated therewith.

A second aspect of the present invention is a method of testing a compound for the ability to affect the rate of dissociation of G_α from $G_{\beta\gamma}$ in a cell. The method comprises: providing a transformed yeast cell as described above; contacting the compound to the cell; and then detecting the rate of dissociation of G_α from $G_{\beta\gamma}$ in the cell. The cells may be provided in an

-3-

aqueous solution, and the contacting step carried out by adding the compound to the aqueous solution.

A third aspect of the present invention is a DNA expression vector capable of expressing a transmembrane protein into the cell membrane of yeast cells. The vector contains a first segment comprising at least a fragment of the extreme amino-terminal coding sequence of a yeast G protein coupled receptor. A second segment is positioned downstream from the first segment (and in correct reading frame therewith), with the second segment comprising a DNA sequence encoding a heterologous G protein coupled receptor.

A fourth aspect of the present invention is a yeast cell transformed by a vector as described above.

15 Brief Description of the Drawings

Figure 1 illustrates the construction of the yeast human β_2 Adrenergic Receptor expression plasmid, pY β AR2.

Figure 2 illustrates h β AR ligand binding to membranes from pY β AR2-transformed yeast cells.

Figure 3 shows a comparison of β -adrenergic agonist effects on pheromone-inducible gene activity. α -MF, 10 μ M α -mating factor; (-) ISO, 50 μ M (-) isoproterenol; (-) ALP, 50 μ M (-) alprenolol; (+) ISO, 100 μ M (+) isoproterenol.

25 Detailed Description of the Invention

Nucleotide bases are abbreviated herein as follows:

30	A=Adenine	G=Guanine
	C=Cytosine	T=Thymine

Amino acid residues are abbreviated herein to either three letters or a single letter as follows:

35	Ala;A=Alanine	Leu;L=Leucine
	Arg;R=Arginine	Lys;K=Lysine
	Asn;N=Asparagine	Met;M=Methionine
	Asp;D=Aspartic acid	Phe;F=Phenylalanine
	Cys;C=Cysteine	Pro;P=Proline

-4-

Gln;Q=Glutamine

Ser;S=Serine

Glu;E=Glutamic acid

Thr;T=Threonine

Gly;G=Glycine

Trp;W=Tryptophan

His;H=Histidine

Tyr;Y=Tyrosine

5 Ile;I=Isoleucine

Val;V=Valine

The term "mammalian" as used herein refers to any mammalian species (e.g., human, mouse, rat, and monkey).

10 The term "heterologous" is used herein with respect to yeast, and hence refers to DNA sequences, proteins, and other materials originating from organisms other than yeast (e.g., mammalian, avian, amphibian), or combinations thereof not naturally found in yeast.

15 The terms "upstream" and "downstream" are used herein to refer to the direction of transcription and translation, with a sequence being transcribed or translated prior to another sequence being referred to as "upstream" of the latter.

20 G proteins are comprised of three subunits: a guanyl-nucleotide binding α subunit; a β subunit; and a γ subunit. G proteins cycle between two forms, depending on whether GDP or GTP is bound thereto. When GDP is bound the G protein exists as an inactive

25 heterotrimer, the $G_{\alpha\beta\gamma}$ complex. When GTP is bound the α subunit dissociates, leaving a $G_{\beta\gamma}$ complex. Importantly, when a $G_{\alpha\beta\gamma}$ complex operatively associates with an activated G protein coupled receptor in a cell membrane, the rate of exchange of GTP for bound GDP is

30 increased and, hence, the rate of dissociation of the bound α subunit from the $G_{\beta\gamma}$ complex increases. This fundamental scheme of events forms the basis for a multiplicity of different cell signaling phenomena. See generally Stryer and Bourne, supra.

35 Any mammalian G protein coupled receptor, and the DNA sequences encoding these receptors, may be employed in practicing the present invention. Examples

-5-

of such receptors include, but are not limited to, dopamine receptors, muscarinic cholinergic receptors, α -adrenergic receptors, β -adrenergic receptors, opiate receptors, cannabinoid receptors, and serotonin receptors. The term receptor as used herein is intended to encompass subtypes of the named receptors, and mutants and homologs thereof, along with the DNA sequences encoding the same.

The human D₁ dopamine receptor cDNA is reported in A. Deary et al., Nature **347**, 72-76 (1990).

The rat D₂ dopamine receptor cDNA is reported in J. Bunzow et al., Nature **336**, 783-787 (1988); see also O. Civelli, et al., PCT Appln. WO 90/05780 (all references cited herein are to be incorporated herein by reference).

Muscarinic cholinergic receptors (various subtypes) are disclosed in E. Peralta et al., Nature **343**, 434 (1988) and K. Fukuda et al., Nature **327**, 623 (1987).

Various subtypes of α_2 -adrenergic receptors are disclosed in J. Regan et al., Proc. Natl. Acad. Sci. USA **85**, 6301 (1988) and in R. Lefkowitz and M. Caron, J. Biol. Chem. **263**, 4993 (1988).

Serotonin receptors (various subtypes) are disclosed in S. Peroutka, Ann. Rev. Neurosci. **11**, 45 (1988).

A cannabinoid receptor is disclosed in L. Matsuda et al., Nature **346**, 561 (1990).

Any DNA sequence which codes for a mammalian G α subunit (G α) may be used to practice the present invention. Examples of mammalian G α subunits include G_s α subunits, G_i α subunits, G_o α subunits, G_z α subunits, and transducin α subunits. See generally Stryer and Bourne, supra. G proteins and subunits useful for practicing the present invention include subtypes, and mutants and homologs thereof, along with the DNA sequences encoding the same.

-6-

Heterologous DNA sequences are expressed in a host by means of an expression vector. An expression vector is a replicable DNA construct in which a DNA sequence encoding the heterologous DNA sequence is operably linked to suitable control sequences capable of effecting the expression of a protein or protein subunit coded for by the heterologous DNA sequence in the intended host. Generally, control sequences include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and (optionally) sequences which control the termination of transcription and translation.

Vectors useful for practicing the present invention include plasmids, viruses (including phage), and integratable DNA fragments (i.e., fragments integratable into the host genome by homologous recombination). The vector may replicate and function independently of the host genome, as in the case of a plasmid, or may integrate into the genome itself, as in the case of an integratable DNA fragment. Suitable vectors will contain replicon and control sequences which are derived from species compatible with the intended expression host. For example, a promoter operable in a host cell is one which binds the RNA polymerase of that cell, and a ribosomal binding site operable in a host cell is one which binds the endogenous ribosomes of that cell.

DNA regions are operably associated when they are functionally related to each other. For example: a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous and, in the case of leader sequences, contiguous and in reading phase.

-7-

Transformed host cells of the present invention are cells which have been transformed or transfected with the vectors constructed using recombinant DNA techniques and express the protein or protein subunit coded for by the heterologous DNA sequences. In general, the host cells are incapable of expressing an endogenous G protein α -subunit (yeast G_α). The host cells do, however, express a complex of the G protein β subunit and the G protein γ subunit ($G_{\beta\gamma}$). The host cells may express endogenous $G_{\beta\gamma}$, or may optionally be engineered to express heterologous $G_{\beta\gamma}$ (e.g., mammalian) in the same manner as they are engineered to express heterologous G_α .

A variety of yeast cultures, and suitable expression vectors for transforming yeast cells, are known. See, e.g., U.S. Patent No. 4,745,057; U.S. Patent No. 4,797,359; U.S. Patent No. 4,615,974; U.S. Patent No. 4,880,734; U.S. Patent No. 4,711,844; and U.S. Patent No. 4,865,989. Saccharomyces cerevisiae is the most commonly used among the yeast, although a number of other strains are commonly available. See, e.g., U.S. Patent No. 4,806,472 (Kluveromyces lactis and expression vectors therefor); 4,855,231 (Pichia pastoris and expression vectors therefor). Yeast vectors may contain an origin of replication from the 2 micron yeast plasmid or an autonomously replicating sequence (ARS), a promoter, DNA encoding the heterologous DNA sequences, sequences for polyadenylation and transcription termination, and a selection gene. An exemplary plasmid is YRp7, (Stinchcomb et al., Nature 282, 39 (1979); Kingsman et al., Gene 7, 141 (1979); Tschemper et al., Gene 10, 157 (1980)). This plasmid contains the TRP1 gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example ATCC No. 44076 or PEP4-1 (Jones, Genetics 85, 12 (1977)). The presence of the TRP1 lesion in the

-8-

yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

Suitable promoting sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., J. Biol. Chem. 255, 2073 (1980) or other glycolytic enzymes (Hess et al., J. Adv. Enzyme Reg. 7, 149 (1968); and Holland et al., Biochemistry 17, 4900 (1978)), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Suitable vectors and promoters for use in yeast expression are further described in R. Hitzeman et al., EPO Publ. No. 73,657. Other promoters, which have the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, and the aforementioned metallothionein and glyceraldehyde-3-phosphate dehydrogenase, as well as enzymes responsible for maltose and galactose utilization.

In constructing suitable expression plasmids, the termination sequences associated with these genes may also be ligated into the expression vector 3' of the heterologous coding sequences to provide polyadenylation and termination of the mRNA.

A novel DNA expression vector described herein which is particularly useful for carrying out the present invention contains a first segment comprising at least a fragment of the extreme amino-terminal coding sequence of a yeast G-protein coupled receptor and a second segment downstream from said

-9-

first segment and in correct reading frame therewith,
the second segment comprising a DNA sequence encoding a
heterologous G protein coupled receptor (e.g., a
mammalian G protein coupled receptor). In a preferred
embodiment, this vector comprises a plasmid. In
constructing such a vector, a fragment of the extreme
amino-terminal coding sequence of the heterologous G
protein coupled receptor may be deleted. The first and
second segments are operatively associated with a
promoter, such as the GAL1 promoter, which is operative
in a yeast cell. Coding sequences for yeast G protein
coupled receptors which may be used in constructing
such vectors are exemplified by the gene sequences
encoding yeast pheromone receptors (e.g., the STE2
gene, which encodes the α -factor receptor, and the STE3
gene, which encodes the α -factor receptor). The levels
of expression obtained from these novel vectors are
enhanced if at least a fragment of the 5'-untranslated
region of a yeast G protein coupled receptor gene
(e.g., a yeast pheromone receptor gene; see above) is
positioned upstream from the first segment and
operatively associated therewith.

Any of a variety of means for detecting the
dissociation of G_α from $G_{\beta\gamma}$ can be used in connection
with the present invention. The cells could be
disrupted and the proportion of these subunits and
complexes determined physically (i.e., by
chromatography). The cells could be disrupted and the
quantity of G_α present assayed directly by assaying for
the enzymatic activity possessed by G_α in isolation
(i.e., the ability to hydrolyze GTP to GDP). Since
whether GTP or GDP is bound to the G protein depends on
whether the G protein exists as a $G_{\beta\gamma}$ or $G_{\alpha\beta\gamma}$ complex,
dissociation can be probed with radiolabelled GTP. As
explained below, morphological changes in the cells can
be observed. A particularly convenient method,
however, is to provide in the cell a third heterologous

-10-

DNA sequence, wherein the third heterologous DNA sequence comprises a pheromone-responsive promoter and an indicator gene positioned downstream from the pheromone-responsive promoter and operatively associated therewith. This sequence can be inserted with a vector, as described in detail herein. With such a sequence in place, the detecting step can be carried out by monitoring the expression of the indicator gene in the cell. Any of a variety of pheromone responsive promoters could be used, examples being the BAR1 gene promoter and the FUS1 gene promoter. Likewise, any of a broad variety of indicator genes could be used, with examples including the HIS3 gene and the LacZ gene.

As noted above, transformed host cells of the present invention express the protein or protein subunit coded for by the heterologous DNA sequence. When expressed, the G protein coupled receptor is located in the host cell membrane (i.e., physically positioned therein in proper orientation for both the stereospecific binding of ligands on the extracellular side of the cell membrane and for functional interaction with G proteins on the cytoplasmic side of the cell membrane).

The ability to control the yeast pheromone response pathway by expression of a heterologous adrenergic receptor and its cognate G protein α -subunit has the potential to facilitate structural and functional characterization of mammalian G protein-coupled receptors. By scoring for responses such as growth arrest or β -galactosidase induction, the functional properties of mutant receptors can now be rapidly tested. Similarly, as additional genes for putative G protein-coupled receptors are isolated, numerous ligands can be screened to identify those with activity toward previously unidentified receptors. See F. Libert *et al.*, Science 244, 569 (1989); M. S. Chee

-11-

et al., Nature 344, 774 (1990). Moreover, as additional genes coding for putative G protein α -subunits are isolated, they can be expressed in cells of the present invention and screened with a variety of G protein coupled receptors and ligands to characterize these subunits. These cells can also be used to screen for compounds which affect receptor-G protein interactions.

Cells of the present invention can be deposited in the wells of microtiter plates in known, predetermined quantities to provide standardized kits useful for screening compounds in accordance with the various screening procedures described above.

The following Examples are provided to further illustrate various aspects of the present invention. They are not to be construed as limiting the invention.

EXAMPLE 1

Construction of the Human β 2-Adrenergic Expression Vector pY β AR2 and Expression in Yeast

To attain high level expression of the human β 2-adrenergic receptor (h β AR) in yeast, a modified h β AR gene was placed under the control of the GAL1 promoter in the multicopy vector, YEp24 (pY β AR2).

Figure 1 illustrates the construction of yeast expression plasmid pY β AR2. In pY β AR2, expression of the h β AR sequence is under the control of the GAL1 promoter. Figure 1A shows the 5'-untranslated region and the first 63 basepairs (bp) of coding sequence of the h β AR gene in pTZNAR, B. O'Dowd et al., J. Biol. Chem. 263, 15985 (1988), which was removed by Aat II cleavage and replaced with a synthetic oligonucleotide corresponding to 11 bp of noncoding and 42 bp of coding sequence from the STE2 gene (SEQ ID NO:1; SEQ ID NO:2). See N. Nakayama et al., EMBO J. 4, 2643 (1985); A. Burkholder and L. Hartwell, Nucleic Acids Res. 13, 8463

-12-

(1985). The resulting plasmid, pTZYNAR, contains the modified h β AR gene flanked by Hind III sites in noncoding sequences with the 3' Hind III site given as SEQ ID NO:3 herein. The Hind III-Hind III fragment was isolated from pTZYNAR and inserted into pAAH5 such that the 3'- untranslated sequence of the modified h β AR gene was followed by 450 bp containing termination sequences from the yeast ADH1 gene. See G. Ammerer, Methods. Enzymol. 101, 192 (1983).

As illustrated in Figure 1B, py β AR2 was constructed by inserting the Bam HI - Bam HI fragment containing h β AR and ADH1 sequences into YEpG24. E. Wyckoff and T. Hsieh, Proc. Natl. Acad. Sci. U.S.A. 85, 6272 (1988). Where maximum expression was sought, cells were cotransformed with plasmid pMTL9 (from Dr. S. Johnston) containing LAC9, a homolog of the S. cerevisiae GAL4 transactivator protein required for GAL1-regulated transcription. J. Salmeron et al., Mol. Cell. Biol. 9, 2950 (1989). Cells grown to late exponential phase were induced in medium containing 3% galactose, supplemented with about 10 μ M alprenolol, and grown for an additional 36 hours. Standard methods for the maintenance of cells were used. See F. Sherman et al., Methods in Yeast Genetics (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1986).

Maximal expression required (i) expression of a transcriptional transactivator protein (LAC9), (ii) replacement of the 5' untranslated and extreme NH₂-terminal coding sequence of the h β AR gene with the corresponding region of the yeast STE2 (α -factor receptor) gene, (iii) induction with galactose when cell growth reached late exponential phase, and, (iv) inclusion of an adrenergic ligand in the growth medium during induction.

The plasmid pY β AR2 was deposited in accordance with the provisions of the Budapest Treaty at the American Type Culture Collection, 12301 Parklawn

-13-

Drive, Rockville, MD 20852 USA, on September 11, 1990,
and has been assigned ATCC Accession No. 40891.

EXAMPLE 2

Binding Affinity of h β AR Ligands in Yeast Transformed with pY β AR2

5 A primary function of cell surface receptors
is to recognize only appropriate ligands among other
extracellular stimuli. Accordingly, ligand binding
affinities were determined to establish the functional
10 integrity of h β AR expressed in yeast. As discussed in
detail below, an antagonist, 125 I-labeled cyanopindolol
(125 I-CYP), bound in a saturable manner and with high
affinity to membranes prepared from pY β AR2-transformed
yeast cells. By displacement of 125 I-CYP with a series
15 of agonists, the order of potency and stereospecificity
expected for h β AR was observed.

SC261 cells (MATa ura3-52 trp1 leu2 prb1-1122
pep4-3 prc1-407) (from Dr. S. Johnston) harboring
pY β AR2 (URA3) and pMTL9 (LEU2) were grown in minimal
20 glucose-free selective media to late log phase (OD₆₀₀ =
5.0), and then induced with the addition of 3%
galactose and 40 μ M alprenolol. After 36 hours, cells
were harvested and spheroplasts were prepared as
described. See E. Wyckoff and T. Hsieh, Proc. Natl.
25 Acad. Sci. U.S.A. 85, 6272 (1988). Briefly, the
spheroplasts were resuspended in 50 mM Tris-HCl pH 7.4,
5 mM EDTA and were lysed by vortex mixing with glass
beads for three one-min periods at 4°C. Crude membranes
were prepared from the lysates and binding assays with
30 125 I-CYP were performed by methods described previously.
See H. Dohlman et al., Biochemistry 29, 2335 (1990).

Figure 2 illustrates h β AR ligand binding to
membranes from pY β AR2-transformed yeast cells. (A) B_{max}
(maximum ligand bound) and K_d (ligand dissociation
35 constant) values were determined by varying 125 I-CYP
concentrations (5 - 400 pM). Specific binding was

-14-

defined as the amount of total binding (circles) minus nonspecific binding measured in the presence of 10 μ M (-) alprenolol (squares). A K_d of 93 pM for 125I-CYP binding was obtained and used to calculate agonist
5 affinities (below). (B) Displacement of 18 pM 125I-CYP with various concentrations of agonists was used to determine apparent low affinity K_i values (non G protein coupled, determined in the presence of 50 μ M GTP) for receptor binding, squares; (-) isoproterenol, circles;
10 (-) epinephrine, downward-pointing triangles; (+) isoproterenol, upward pointing triangles; (-) norepinephrine.

COMPARATIVE EXAMPLE A

Ligand Binding Affinity for h β AR Expressed 15 in Yeast and Mammalian Cells

The binding data of Figures 2 (A) and (B) were analyzed by nonlinear least squares regression, see A. DeLean et al., Mol. Pharmacol. 21, (1982), and are presented in Table I. Values given are averages of
20 measurements in triplicate, and are representative of 2 - 3 experiments. Binding affinities in yeast were nearly identical to those observed previously for h β AR expressed in mammalian cells.

-15-

Tabl 1

Comparison of ligand Binding Parameters for High Level Expression of Human β -Adrenergic Receptor in Yeast and COS-7 Cells*

5

	Yeast	Monkey	
	SC261 (pYβAR2, pMTL9)	COS-7 (pBC12:hβAR)	

	125 _I -CYP:		
10	¹ K _d	0.093 nM ± 0.013	0.110 nM ± 0.009
	² B _{max}	115 pmol/mg	24 pmol/mg

	³ K _i (M):		
	(-) isoproterenol	103 ± 26	130 ± 15
15	(+) isoproterenol	3670 ± 420	4000 ± 184
	(-) epinephrine	664 ± 123	360 ± 30
	(-) norepinephrine	6000 ± 1383	5800 ± 373

20	*Values derived from Fig. 2 and H. Dohlman <u>et al.</u> , <u>Biochemistry</u> 29, 2335 (1990).; ± S.E.		
	¹ K _d , ligand dissociation constant		
	² B _{max} , maximum ligand bound		
	³ K _i , inhibition constant		

EXAMPLE 3

25 Agonist-Dependent Activation of Mating Signal Transduction in Yeast Expressing h β AR

A second major function of a receptor is agonist-dependent regulation of downstream components in the signal transduction pathway. Because the pheromone-responsive effector in yeast is not known, indirect biological assays are the most useful indicators of receptor functionality. See K. Blumer and J. Th rner, Annu.Rev. Physiol. in press; I. Herskowitz, Microbiol. Rev. 52, 536 (1988). In yeast

30

-16-

cells expressing high concentrations of h β AR, no agonist-dependent activation of the mating signal transduction pathway could be detected by any of the typical in vivo assays; for example, imposition of G1 arrest, induction of gene expression, alteration of morphology (so-called "shmoo" formation) or stimulation of mating. A likely explanation for the absence of responsiveness is that h β AR was unable to couple with the endogenous yeast G protein.

10

EXAMPLE 4

Coexpression of h β AR and Mammalian G_s α -Subunit in Yeast

Expression of a mammalian G_s α -subunit can correct the growth defect in yeast cells lacking the corresponding endogenous protein encoded by the GPA1 gene. See C. Dietzel and J. Kurjan, Cell **50**, 1001 (1987). Moreover, specificity of receptor coupling in mammalian cells is conferred by the α -subunit of G proteins. See L. Stryer and H. Bourne, Annu. Rev. Cell Biol. **2**, 391 (1988). Thus, coexpression of h β AR and a mammalian G_s α -subunit (G_s α) in yeast was attempted to render the yeast responsive to adrenergic ligands. Accordingly, a cDNA encoding rat G_s α under the control of the copper-inducible CUP1 promoter was introduced on a second plasmid, pYSK136G α s. See C. Dietzel and J. Kurjan, Cell **50**, 1001 (1987). In yeast (NNY19) coexpressing h β AR and rat G_s α , but containing wild-type GPA1, no adrenergic agonist-induced shmoo formation, a characteristic morphological change of yeast in response to mating pheromone, was observed.

30

EXAMPLE 5

Coexpression of h β AR and Mammalian G_s α -Subunit in Yeast Lacking an Endogenous G Protein α -Subunit

To prevent interference by the endogenous yeast G protein α -subunit, gpa1 mutant cells (strain

35

-17-

8c) were used.

Yeast strain 8c (MATa ura3 leu2 his3 trp1
gpa1::HIS3), I. Miyajima et al., Cell 50, 1011 (1987),
carrying plasmids pYSK136G α s (TRP1), C. Dietzel and J.
Kurjan, Cell 50, 1001 (1987), pMTL9 (LEU2), J. Salmeron
et al., Mol. Cell. Biol. 9, 2950 (1989), and pY β AR2
(URA3) was maintained on glucose-free minimal selective
plates containing 3% glycerol, 2% lactic acid, 50 μ M
CuSO₄ and 3% galactose. Colonies were transferred to
similar plates containing 0.5 mM ascorbic acid and the
indicated adrenergic ligand(s). After 16-20 hours at
30°C, the colonies were transferred to similar liquid
media at a density of 10⁶-10⁷ cells/ml and examined by
phase contrast microscopy.

Morphologies of yeast cells cotransformed
with pY β AR2, pMTL9, and pYSK136G α s were examined after
incubation with (A) no adrenergic agent; (B) 100 μ M (-)
isoproterenol; (C) 100 μ M (-) isoproterenol and 50 μ M
(-) alprenolol; and (D) 100 μ M (+) isoproterenol.
Results showed that treatment of 8c cells coexpressing
h β AR and rat G α s with the β -adrenergic agonist
isoproterenol indeed induced shmoo formation, and that
this effect was blocked by the specific antagonist
alprenolol.

25

EXAMPLE 6

Coexpression of h β AR and Mammalian G α -Subunit in Yeast Containing a β -Galactosidase Signal Sequence

The isoproterenol-induced morphological
response of 8c cells coexpressing h β AR and rat G α s
suggested that these components can couple to each
other and to downstream components of the pheromone
response pathway in yeast lacking the endogenous
G α -subunit. To confirm that the pheromone signaling
pathway was activated by h β AR and rat G α s, agonist
induction of the pheromone-responsive FUS1 gene
promoter was measured in a strain of yeast derived from

-18-

8c cells (8c1) in which a FUS1-lacZ gene fusion had been stably integrated into the genome. See S. Nomoto et al., EMBO J. 9, 691 (1990).

Strains 8c (Fig. 3, legend) and NNY19 (MATa ura3 leu2 his3 trp1 lys2 FUS1-LacZ::LEU2) were modified by integrative transformation with YIpFUS102 (LEU2), S. Nomoto et al., supra, and designated 8c1 and NNY19, respectively. These strains were transformed with pY β AR2 and pYSK136G α s and maintained on minimal selective plates containing glucose and 50 μ M CuSO₄. Colonies were inoculated into minimal selective media (3% glycerol, 2% lactic acid, 50 μ M CuSO₄), grown to early log phase (OD₆₀₀ = 1.0), and induced for 12 hours by addition of 3% galactose. Cells were washed and resuspended in induction media (OD₆₀₀ = 5.0) containing 0.5 mM ascorbic acid and the indicated ligands. After a 4 hour incubation at 30°C, cells were harvested, resuspended into 1 ml of Z-buffer, see J. Miller, Experiments in Molecular Genetics (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1972), supplemented with 0.0075% SDS, and β -galactosidase activities were determined in 3 - 4 independent experiments as described previously. See J. Miller, supra.

Figure 3 shows a comparison of β -adrenergic agonist effects on pheromone-inducible gene activity. α -MF, 10 μ M α -mating factor; (-) ISO, 50 μ M (-) isoproterenol; (-) ALP, 50 μ M (-) alprenolol; (+) ISO, 100 μ M (+) isoproterenol. In 8c1 (gpa1) cells coexpressing h β AR and rat G_s α , a dramatic isoproterenol-stimulated induction of β -galactosidase activity was observed. Agonist stimulation was stereoselective and was blocked by addition of a specific antagonist. Agonist responsiveness was dependent on expression of both h β AR and rat G_s α , and required a strain in which the endogenous G protein α -subunit was disrupted. The final β -galactosidase activity achieved in response to isopr terenol in transformed 8c1 cells was comparable

-19-

t that induced by α -factor in nontransformed cells that express GPA1 (NNY19), although basal β -galactosidase activity in NNY19 cells was considerably lower than in 8c1 cells. Taken together, 5 our results indicated that coexpression of h β AR and rat $G_s\alpha$ was sufficient to place under catecholamine control key aspects of the mating signal transduction pathway in yeast. However, the adrenergic agonist did not stimulate mating in either 8c cells or NNY19 cells 10 coexpressing h β AR and rat $G_s\alpha$, in agreement with recent observations that yeast pheromone receptors, in addition to binding pheromones, participate in other recognition events required for mating. See A. Bender and G. Sprague, Genetics 121, 463 (1989).

15 h β AR stimulates adenylate cyclase in animal cells via the action of the α -subunit of its G protein. In contrast, mating factor receptors in yeast trigger their effector via the action of the $\beta\gamma$ subunits. M. Whiteway et al., Cell 56, 476 (1989). Our present 20 results indicate that activation of h β AR in yeast leads to dissociation of mammalian $G_s\alpha$ from yeast $\beta\gamma$, and it is the $\beta\gamma$ subunits that presumably elicit the response.

The foregoing examples are illustrative of the present invention, and are not to be construed as 25 limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

-20-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: King, Klim
Dohlman, Henrik G.
Caron, Mark G.
Lefkowitz, Robert J.

(ii) TITLE OF INVENTION: Expression of G Protein Coupled
Receptors in Yeast

(iii) NUMBER OF SEQUENCES: 3

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Kenneth D. Sibley; Bell, Seltzer, Park and
Gibson
(B) STREET: Post Office Drawer 34009
(C) CITY: Charlotte
(D) STATE: North Carolina
(E) COUNTRY: U.S.A.
(F) ZIP: 28234

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: PCT/US91/06605
(B) FILING DATE: 12-SEP-1991
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 07/581714
(B) FILING DATE: 13-SEP-1990

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Sibley, Kenneth D.
(B) REGISTRATION NUMBER: 31,665
(C) REFERENCE/DOCKET NUMBER: 5405-17-1

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 919-881-3140
(B) TELEFAX: 919-881-3175
(C) TELEX: 575102

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 80 base pairs

-21-

(B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 30..80

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAATTCAACG	TTGGATCCAA	GAATCAAAA	ATG	TCT	GAT	GCG	GCT	CCT	TCA	TTG	53
			Met	Ser	Asp	Ala	Ala	Pro	Ser	Leu	
			1				5				

AGC	AAT	CTA	TTT	TAT	GAC	GTC	ACG	CAG		80
Ser	Asn	Leu	Phe	Tyr	Asp	Val	Thr	Gln		
10					15					

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Ser	Asp	Ala	Ala	Pro	Ser	Leu	Ser	Asn	Leu	Phe	Tyr	Asp	Val	Thr
1				5				10					15		
															Gln

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AACGTT

6

-22-

THAT WHICH IS CLAIMED IS:

1. A transformed yeast cell containing a first heterologous DNA sequence which codes for a mammalian G protein coupled receptor and a second heterologous DNA sequence which codes for a mammalian G protein α subunit (mammalian G_α), wherein said first and second heterologous DNA sequences are capable of expression in said cell, and wherein said cell is incapable of expressing an endogenous G protein α -subunit (yeast G_α).
5
2. A transformed yeast cell according to claim 1, wherein said first heterologous DNA sequence is carried by a plasmid.
3. A transformed yeast cell according to claim 1, wherein said second heterologous DNA sequence is carried by a plasmid.
4. A transformed yeast cell according to claim 1, wherein said mammalian G protein α subunit is selected from the group consisting of G_s α subunits, G_i α subunits, G_o α subunits, G_z α subunits, and transducin α subunits.
5
5. A transformed yeast cell according to claim 1 which expresses a complex of the G protein β subunit and the G protein γ subunit ($G_{\beta\gamma}$).
6. A transformed yeast cell according to claim 5 which express s ndogenous $G_{\beta\gamma}$.

-23-

7. A transformed yeast cell according to claim 1, wherein said first heterologous DNA sequence codes for a mammalian G protein-coupled receptor selected from the group consisting of dopamine
5 receptors, muscarinic cholinergic receptors, α -adrenergic receptors, β -adrenergic receptors, opiate receptors, cannabinoid receptors, and serotonin receptors.

8. A transformed yeast cell according to claim 1 further comprising a third heterologous DNA sequence, wherein said third heterologous DNA sequence comprises a pheromone-responsive promoter and an
5 indicator gene positioned downstream from said pheromone-responsive promoter and operatively associated therewith.

9. A transformed yeast cell according to claim 8, wherein said pheromone responsive promoter is selected from the group consisting of the BAR1 gene promoter and the FUS1 gene promoter, and wherein said
5 indicator gene is selected from the group consisting of the HIS3 gene and the LacZ gene.

-24-

10. A method of testing a compound for the ability to affect the rate of dissociation of G_α from $G_{\beta\gamma}$ in a cell, comprising:

- 5 providing a transformed yeast cell containing a first heterologous DNA sequence which codes for a mammalian G protein coupled receptor and a second heterologous DNA sequence which codes for a mammalian G_α , wherein said first and second heterologous DNA sequences are capable of expression in said cell,
- 10 wherein said cell is incapable of expressing endogenous G_α , and wherein said cell expresses $G_{\beta\gamma}$;
- contacting said compound to said cell; and
- detecting the rate of dissociation of G_α from $G_{\beta\gamma}$ in said cell.

11. A method according to claim 10, wherein said yeast cells are provided in an aqueous solution and said contacting step is carried out by adding said compound to said aqueous solution.

12. A method according to claim 10, wherein said mammalian G protein α subunit is selected from the group consisting of G_s α subunits, G_i α subunits, G_o α subunits, G_z α subunits, and transducin α subunits.

13. A method according to claim 10, wherein said yeast cell expresses endogenous $G_{\beta\gamma}$.

-25-

14. A method according to claim 10, wherein
said first heterologous DNA sequence codes for a
mammalian G protein-coupled receptor selected from the
group consisting of dopamine receptors, muscarinic
5 cholinergic receptors, α -adrenergic receptors, β -
adrenergic receptors, opiate receptors, cannabinoid
receptors, and serotonin receptors.

15. A method according to claim 10, said
yeast cell further comprising a third heterologous DNA
sequence, wherein said third heterologous DNA sequence
comprises a pheromone-responsive promoter and an
5 indicator gene positioned downstream from said
pheromone-responsive promoter and operatively
associated therewith;

and wherein said detecting step is carried
out by monitoring the expression of said indicator gene
10 in said cell.

16. A DNA expression vector capable of
expressing a transmembrane protein into the cell
membrane of yeast cells, comprising:

5 a first segment comprising at least a
fragment of the extreme amino-terminal coding sequence
of a yeast G protein coupled receptor; and

a second segment downstream from said first
segment and in correct reading frame therewith, said
second segment comprising a DNA sequence encoding a
10 heterologous G protein coupled receptor.

17. A DNA expression vector according to
claim 16, wherein a fragment of the extreme amino-
terminal coding sequence of said heterologous G protein
coupled receptor is absent.

-26-

18. A DNA expression vector according to claim 16, wherein said first and second segments are operatively associated with a promoter operative in a yeast cell.

19. A DNA expression vector according to claim 18, wherein said promoter is the GAL1 promoter.

20. A DNA expression vector according to claim 16, wherein said first segment comprises at least a fragment of the extreme amino-terminal coding sequence of a yeast pheromone receptor.

21. A DNA expression vector according to claim 16, wherein said first segment comprises at least a fragment of the extreme amino-terminal coding sequence of a yeast pheromone receptor selected from the group consisting of the STE2 gene and the STE3 gene.

22. A DNA expression vector according to claim 16, further comprising at least a fragment of the 5'-untranslated region of a yeast G protein coupled receptor gene positioned upstream from said first segment and operatively associated therewith.

-27-

23. A DNA expression vector according to claim 16, further comprising at least a fragment of the 5'-untranslated region of a yeast pheromone receptor gene positioned upstream from said first segment and operatively associated therewith.

24. A DNA expression vector according to claim 23, wherein said yeast pheromone receptor gene is selected from the group consisting of the STE2 gene and the STE3 gene.


25. A DNA expression vector according to claim 16, said vector comprising a plasmid.

26. A DNA expression vector according to claim 16, said second segment comprising a DNA sequence encoding a mammalian G protein coupled receptor.

27. A DNA expression vector according to claim 16, said second segment comprising a DNA sequence encoding a mammalian G protein-coupled receptor selected from the group consisting of dopamine receptors, muscarinic cholinergic receptors, α -adrenergic receptors, β -adrenergic receptors, opiate receptors, cannabinoid receptors, and serotonin receptors.

28. A yeast cell carrying a DNA expression vector according to claim 16.

FIG. 1A.



1/3







FIG. 1B.

FIG. 2A.FIG. 2B.

FIG. 3.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/06605

I. CLASSIFICATION AND SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶
 According to International Patent Classification (IPC) or to both National Classification and IPC
 IPC(5): C12N 1/20; C12! 1/00; G01N 33/48, 33/566
 USCL.: 435/4,252.3; 436/63,501

II. FIELDS SEARCHED
 Minimum Documentation Searched ⁷

Classification System	Classification Symbols
US.CL.	435/4, 252.3; 436/63,501

 Documentation Searched other than Minimum Documentation
 to the Extent that such Documents are Included in the Fields Searched ⁸

III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	Annual Review of Cell Biology, Volume 2, issued 1986. Stryer et al.. "G Proteins: A Family of Signal Transducers," pages 391-419. see entire document.	1-28
Y	WO.A. 90/05780 (Civelli et al.) 31 May 1990. see column 29. line 26-column 30. line 8.	1-28

- * Special categories of cited documents: ¹⁰
- "A" document defining the general state of the art which is not considered to be of particular relevance
 - "E" earlier document but published on or after the international filing date
 - "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 - "O" document referring to an oral disclosure, use, exhibition or other means
 - "P" document published prior to the international filing date but later than the priority date claimed
 - "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 - "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
 - "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 - "&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

01 November 1991

International Searching Authority

ISA/US

Date of Mailing of this International Search Report

18 DEC 1991

Signature of Authorized Officer

Lori Yuan